

Your name: \_\_\_\_\_ Date of birth: \_\_\_\_/\_\_\_\_/\_\_\_\_ Today's date: \_\_\_\_/\_\_\_\_/\_\_\_\_  
(mo.) (day) (yr.) (mo.) (day) (yr.)



## Do I Need Any Vaccinations Today?

Many adults are behind on their vaccinations. This questionnaire will help you and your healthcare provider determine if you need any vaccinations today. Please check the boxes that apply to you.

### Influenza vaccination

- I haven't had my annual influenza vaccination yet this season — so I need it now.

### Pneumococcal vaccination (PPSV23, PCV13)

- I am age 65 or older, and I have never had a pneumococcal shot.
- I am age 65 or older and have had 1 or 2 doses of pneumococcal vaccine when I was younger than age 65; it has been 5 years or more since the last shot.
- I am younger than age 65, I have not been vaccinated against pneumococcal disease, and at least one of the following applies to me:
- I smoke cigarettes.
  - I have heart, lung (including asthma), liver, kidney, or sickle cell disease; diabetes; or alcoholism.
- I am younger than age 65, I have or have not been vaccinated against pneumococcal disease, and at least one of the following applies to me:
- I have a weakened immune system due to cancer, Hodgkin's disease, leukemia, lymphoma, multiple myeloma, kidney failure, HIV/AIDS; or I am receiving radiation therapy; or I am on medication that suppresses my immune system.
  - I have had an organ or bone marrow transplant.
  - I have had my spleen removed, have had or will have a cochlear implant, or have leaking spinal fluid.
- I live in a nursing home or other long-term care facility, and I have never had a pneumococcal shot.

### Tetanus-, diphtheria-, and pertussis (whooping cough)-containing vaccination (e.g., DTP, DTaP, Tdap, or Td)

- I have not had a pertussis-containing vaccine (Tdap) as an adolescent or adult (this applies to pregnant women, too).
- I have not yet had at least 3 tetanus- and diphtheria-containing shots.
- I have had at least 3 tetanus- and diphtheria-containing shots in my lifetime, but I believe it's been 10 years or more since I received my last shot.
- I have no idea if I ever received any tetanus- and diphtheria-containing shots in school, the military, or elsewhere.

### Measles-Mumps-Rubella (MMR) vaccination

- I was born in 1957 or later and never received an MMR shot or I don't know if I received a shot.
- I am a woman thinking about a future pregnancy and do not know if I'm immune to rubella.
- I am a healthcare worker, I do not have a history of measles or mumps, and I've had only one dose of MMR vaccine.
- I was born in 1957 or later, and I am included in one of the following groups for whom 2 MMR shots are recommended, but I have received only 1 shot.
- I am entering college or a post-high school educational institution.
  - I had a blood test that shows I do not have immunity to measles, mumps, or rubella.
  - I am planning to travel internationally.

### Human papillomavirus (HPV) vaccination

- I am a woman age 26 or younger and haven't completed a series of shots against human papillomavirus.
- I am a man age 21 or younger and haven't completed a 3-dose series of shots against human papillomavirus.
- I am a man age 22 through 26, I have not completed a 3-dose series of HPV vaccine, and at least 1 of the following applies to me:
- I want to be protected from HPV.
  - I have a weakened immune system as a result of infection (including HIV), disease, or medications.
  - I have sex with men.
- I am older than age 26 and although I started the series when I was younger, I never completed it.

(continued on page 2)

**Hepatitis A vaccination**

- I want to be vaccinated to avoid getting hepatitis A and spreading it to others.
- I was vaccinated with hepatitis A vaccine in the past but never received the second shot.
- I might have been exposed to the hepatitis A virus in the past 2 weeks.
- I am in one of the following risk groups, and I haven't completed the 2-dose series of hepatitis A shots:
  - I travel or plan to travel in countries where hepatitis A is common.<sup>1,2</sup>
  - I use street drugs.
  - I have (or will have) contact with an adopted child within the first 60 days of their arrival from a country where hepatitis A is common.<sup>2</sup>
  - I have chronic liver disease.
  - I am a man who has sex with men.
  - I have a clotting factor disorder.
  - I work with HAV-infected primates or with HAV in a research laboratory setting.

**Hepatitis B vaccination**

- I want to be vaccinated to avoid getting hepatitis B and spreading it to others.
- I am age 18 or younger and haven't completed the series of hepatitis B shots.
- I was vaccinated with hepatitis B vaccine in the past but never completed the full 3-dose series.
- I am in one of the following risk groups, and I haven't completed the series of hepatitis B shots:
  - I am sexually active and am not in a long-term, mutually monogamous relationship.
  - I inject street drugs.
  - I am a man who has sex with men.
  - I have chronic liver disease.
  - I am an immigrant, or my parents are immigrants, from an area of the world where hepatitis B is common, so I need testing and may need vaccination.<sup>3,4</sup>
  - I am or will be on kidney dialysis.
  - I live with or am a sex partner of a person with hepatitis B.
  - I am younger than age 60 years and have diabetes.
  - I have been diagnosed with a sexually transmitted disease.
  - I am a healthcare or public safety worker who is exposed to blood or other body fluids.
  - I have been diagnosed with HIV.
  - I provide direct services for people with developmental disabilities.
  - I travel or plan to travel outside the U.S.<sup>1,3</sup>

**Chickenpox (varicella) vaccination**

- I was born in 1980 or later and have never had chickenpox or the vaccine, or I just don't know.
- I was born before 1980 and am either a healthcare worker or foreign born, and am not sure if I've had chickenpox or not.
- I have received 1 dose of varicella vaccine in the past, but never got a second shot.

**Meningococcal vaccination**

- I am age 18 or younger and haven't received a meningococcal shot.
- I am age 21 or younger and am (or will be) in college, living in a residence hall, and haven't had a meningococcal shot since my 16th birthday.
- I am traveling to an area of the world where meningococcal disease is common.<sup>1</sup>
- I have sickle cell disease, or my spleen isn't working or has been removed, or I have a persistent complement component deficiency.
- I am a microbiologist routinely exposed to isolates of *Neisseria meningitidis*.
- I was previously vaccinated 5 or more years ago and continue to be at risk for meningococcal disease. Note: this does not apply to students whose only risk factor is attending college.

**Shingles (zoster) vaccination**

- I am an adult age 60 or older and haven't had a shingles shot.

**Note:** Adults who travel may need additional vaccinations, such as polio or others. Talk to your healthcare provider.

**Footnotes**

1. Call your local travel clinic to find out if additional vaccines are recommended.
2. Countries where hepatitis A is common include all countries other than the U.S., Western Europe, Canada, Japan, Australia, and New Zealand.
3. Areas with high rates of hepatitis B include Africa, China, Korea, Southeast Asia including Indonesia and the Philippines, South and Western Pacific Islands, interior Amazon Basin, certain parts of the Caribbean (i.e., Haiti and the Dominican Republic), and the Middle East except Israel. Areas with moderate rates include South Central and Southwest Asia, Israel, Japan, Eastern and Southern Europe, Russia, and most of Central and South America.
4. Most adults from moderate- or high-risk areas of the world do not know their hepatitis B status. All patients from these areas need hepatitis B blood tests to determine if they have been previously infected. The first hepatitis B shot can be given during the same visit as the blood tests but only after the blood is drawn.